



Starfish smooth muscle relaxing activity of SALMFamide isotype peptide and its analog derived from starfish, *Asterias rubens*

Anastasia Kubarova, Hye-Jin Go, Hye Young Oh, Nam Gyu Park*

Department of Biotechnology, College of Fisheries Sciences, Pukyong National University, Busan 48513, Korea

Abstract

An organism's physiological processes and behaviors are regulated by neuropeptides and hormone peptides. The first neuropeptide identified from echinoderms is SALMFamide. The two most well-studied SALMFamide neuropeptides are S1 and S2, which possess myoactivity on apical muscle, tube feet, and the cardiac stomach of starfishes. However, neuropeptide candidates identified from SALMFamide's precursor protein sequence have not been investigated. This study aims to compare the bioactivity of SALMFamide neuropeptides from the starfish *Asterias rubens* using various starfish muscle preparations. In this study, the bioactivity of the L-type SALMFamide neuropeptides from the starfish *A. rubens*, AYHTGLPFamide (SALMFa-A) and the derivative AYHSALMFamide (SALMFa-B) was investigated. The neuropeptides were applied on *Asterias amurensis* apical muscle, tube feet, which revealed that the neuropeptides exhibit relaxing activity on apical muscle but no activity on tube feet. The native SALMFa-A peptide had lower relaxing activity on the apical muscle compared to the derivative peptide SALMFa-B. The relaxing activity of two neuropeptides also was compared with those on the apical muscle of *Patiria pectinifera*, which revealed relaxing activity as well as SALMFamide-S1 and S2 neuropeptides. Moreover, the investigation of SALMFa-A and SALMFa-B peptides' bioactivity on *P. pectinifera* cardiac stomach muscle also showed slight relaxing activity.

Keywords: *Asterias rubens*, *Asterias amurensis*, Myoactivity, Neuropeptide, *Patiria pectinifera*, Starfish

Introduction

Neuropeptides are molecules comprising 5 to 50 amino acids that promote intercellular communication (Vishwanatha et al., 2017). Neuropeptides promote the connection between the nervous and endocrine systems (Klavdieva, 1995). The physi-

ological processes and behaviors of starfishes such as locomotion, feeding, and reproduction are controlled and regulated by neuropeptides or hormone peptides (Birenheide et al., 1998; Byrne et al., 2019; Kato et al., 2009; Thorndyke & Carnevali, 2001; Tinoco et al., 2021). To date, the several muscle relaxant and contractile neuropeptides were isolated or identified within

Received: Sep 7, 2022 Revised: Oct 20, 2022 Accepted: Oct 28, 2022

*Corresponding author: Nam Gyu Park

Department of Biotechnology, College of Fisheries Sciences, Pukyong National University, Busan 48513, Korea

Tel: +82-51-629-5867, E-mail: ngpark@pknu.ac.kr

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © 2022 The Korean Society of Fisheries and Aquatic Science

the echinoderm phylum (Kim et al., 2016; Semmens & Elphick, 2017; Semmens et al., 2013).

One of the most studied neuropeptides from echinoderms are SALMFamide peptides, which are present in starfish, sea urchin, and sea cucumber (Elphick, 2014; Elphick et al., 1991). The SALMFamide peptides were purified from *Asterias rubens* and *Asterias forbesi* using FMRFamide antibody (Elphick et al., 1989). SALMFamide is classified into two types: L-type and F-type. The L-type has a pentapeptide motif of SxLxF at the C-terminus, and the F-type has an SxFxF motif at the C-terminus (Elphick et al., 2013). The most studied L-type of SALMFamides are two peptides S1 and S2. SALMFamide-1 (SALMFa-S1) peptide is an octapeptide composed of 8 amino acids, and the amino acid sequence is GFNSALMF. SALMFamide-2 (SALMFa-S2) peptide is a dodecapeptide consisting of 12 amino acids, and the amino acid sequence is SGPYSFNSGLTF. SALMFamide-S1 and SALMFamide-S2 cause the relaxing effect on tube feet, cardiac stomach, and induce the inversion of the cardiac stomach in starfish (Elphick et al., 1991; Elphick et al., 1995; Melarange & Elphick, 2003). SALMFamide-S1 was proposed to take part in promoting growth during regeneration (Thorndyke & Carnevali, 2001). In a recent study, SALMFamide-S1 transcriptional expression during the regeneration process was reported (Byrne et al., 2019). The precursor protein of *A. rubens* SALMFamide comprises additional putative peptides that have different sequences as shown in Fig. 1 (GenBank: ALJ99974.1). Studies on the bioactivity of these putative peptides have not yet been conducted.

The L-type SALMFamide precursor along with SALMFamide-S1 comprises additional 6 putative peptides, which contain SALMFamide and GLPFamide motifs. The two putative

peptides with similar N-terminal motifs and different C-terminal motif were selected for this study. The native AYHTGLPFamide peptide was indicated as SALMFamide-A (SALMFa-A) (Fig. 1). To investigate the effect of C-terminal SALMFamide motif on bioactivity, the proline of the putative AYHSALPFamide peptide was substituted with methionine AYHSALMFamide. This derivative AYHSALMFamide peptide was indicated as SALMFamide-B (SALMFa-B). The SALMFamide-S1 and SALMFamide-S2 neuropeptides were used as control peptides (Fig. 1).

Materials and Methods

Animals

The starfishes *Patiria pectinifera* and *Asterias amurensis* were collected in Busan, Korea. This study did not require local agency/ethics committee approval as experimental work on starfish is not regulated.

Peptides synthesis

The four neuropeptides used in this study were synthesized and purchased from Pepton (Daejeon, Korea).

Bioactivity measurement

To measure synthetic peptides' *in vitro* activities and pharmacology, three muscle preparation of muscle tissue from the starfishes *P. pectinifera* and *A. amurensis* were used (Elphick et al., 1995). All measurements were performed in 55 mM Mg²⁺ artificial seawater (ASW) with the following compounds: (mM) NaCl 445, KCl 10, CaCl₂·2H₂O 10, MgCl₂·6H₂O 55, Glucose 10, Tris-HCl 10 (pH 7.8). The muscle preparations were connected to a physiography system equipped with a force-displacement




Fig. 1. L-type SALMFamide precursor (*Asterias rubens*). The precursor protein comprises 6 predicted isotypes. Peptides used in this study are shown in red, putative peptides are shown in blue, dibasic cleavage sites (KR) are shown in green.

transducer (Type 45196A; NEC-Sanei Instrument, Tokyo, Japan). The relaxing activity of muscle was monitored and registered by a recorder (WR7300; GRAPHTEC, Yokohama, Japan) through an amplifier (AS1302; NEC-Sanei Instrument).

Prior to testing the SALMFamide peptides' relaxing activity, the apical muscle was precontracted by acetylcholine (Ach) 5×10^{-6} M while tube feet and cardiac stomach were activated by carbachol (Carb) 5×10^{-6} M. To measure the pharmacological values, synthetic peptides were dissolved with a calculated amount of distilled water to concentration of 1×10^{-2} M and were logarithmically diluted. Then 20 μ L of peptides were applied in the 2 mL polypropylene chamber to obtain the proper final concentration. The activity was observed and measured for 2 min after the application. After each application muscle was washed by ASW buffer and stabilized for 15 min prior to the next application.

Muscle preparations

Apical muscle

The muscle preparation starts with the starfish dissection by the removal of starfish eyespot followed by the separation of the aboral and oral side using scissors. To prepare the starfish apical muscle, the pyloric caeca were removed from the center of arms. Then each arm was cut, and the apical muscle was removed from the inner side of the body wall. Lastly, approximately 10 mm of apical muscle was cut, and both ends were tied with cotton threads. The muscle preparation was vertically fixed in a 2 mL polypropylene chamber with aeration, where the bottom end is connected to a silver wire on the bottom of the chamber, and the top end is connected to a physiography system.

Tube feet

To prepare the starfish tube feet, the ambulacral plate was cut using scalpel. Then, the tube feet were cut together with an ambulacral ossicle and ampulla. Both ends of the muscle were tied with cotton threads, where on the bottom side, the loop was previously made. The top end was tied with a tube feet sucker, whereas the bottom side was tied with an ambulacral ossicle. And finally, the tube feet preparation was vertically fixed in a 2 mL polypropylene chamber in the same manner with apical muscle.

Cardiac stomach

To prepare the cardiac stomach muscle, the extrinsic retractor strands that are attached the cardiac stomach to the ambula-

crum were cut carefully to separate the stomach from the inner side. Both ends of the muscle were tied with cotton threads. The bottom end of the loop was tied with the aboral part whereas the top end was tied with the oral opening part of the cardiac stomach. Finally, the muscle preparation was vertically fixed in a 2 mL polypropylene chamber in the same manner with previous muscles.

Statistical analysis

For the statistical analyses, one-way analyses of variance (ANOVA) followed by the Bonferroni multiple comparisons test were performed by GraphPad Prism 8.4 for Windows (GraphPad Software, CA, USA). The means denoted with different letters at the top of the columns indicate statistically significant differences ($p < 0.05$). To determine the pharmacological properties the sigmoidal-shaped four-parameter dose-response curves were constructed using GraphPad Prism 8.4. The peptide efficacy (E_{max}) is based on the best-fit top value on a dose-dependent curve constructed by the *in vitro* bioassay values, which is calculated as a relative percentage of the muscle maximum contraction induced by Ach or Carb. All values are shown with mean SD. The statistically significant differences ($p < 0.05$) were determined by one-way ANOVA or two-way ANOVA followed by the Bonferroni multiple range test using GraphPad Prism 8.4.

Results

The pharmacological activity of SALMFamide peptides and a derivative was investigated. The investigation conducted using apical muscle of *A. amurensis* revealed comparable relaxing activity of SALMFa-A and SALMFa-B (Fig. 2A). However, SALMFa-A and SALMFa-B showed lower relaxing activity compared to the control SALMFa-S1 and SALMFa-S2 peptides (Fig. 2B). The weakest relaxing activity was exhibited by the native SALMFa-A peptide (Fig. 2C). The results revealed following E_{max} values: the SALMFa-S1, SALMFa-S2, SALMFa-A and SALMFa-B had E_{max} (%) values of $19.92 \pm 0.99\%$, $22.10 \pm 2.77\%$, $11.41 \pm 0.61\%$ and $15.32 \pm 0.98\%$, respectively. The order of relaxing effect on *A. amurensis* apical muscle was: SALMFa-S2 > SALMFa-S1 > SALMFa-B > SALMFa-A. The threshold concentration (TC) relaxing response for SALMFa-S1, SALMFa-A, and SALMFa-B were observed at 10^{-8} M. Meanwhile, the TC of SALMFa-S2 was observed at 5×10^{-8} M concentration (Table 1).

The investigation of SALMFa-A and SALMFa-B peptides' relaxing activity, performed on apical muscle of *P. pectinifera*,




Fig. 2. Investigation of relaxing effects of SALMFamide peptides on *Asterias amurensis* apical muscle. (A) Illustration of each peptide's relaxing responses on apical muscle of *A. amurensis*. The muscle precontractions were caused by Ach, 5×10^{-6} M. (B) Concentration dependent response curve of SALMFamide peptides on *A. amurensis* apical muscle preparation. The sigmoidal-shaped four-parameter dose-response curves were constructed using GraphPad PRISM 8.4. The relaxing response expressed as % of the maximal response to Ach, 5×10^{-6} M. Data represent the mean \pm SD determined from three separate experiments ($n = 3$). (C) Comparison of the SALMFamide peptides relaxing activity efficacy (E_{max} value). The asterisks on the top of bars denote statistically significant differences ($p < 0.05$) determined by one-way analyses of variance followed by the Bonferroni multiple range test. Ach, acetylcholine.

Table 1. Pharmacological activities of SALMFamide peptides on the apical muscle of starfishes *Asterias amurensis* and *Patiria pectinifera*

Tissue	<i>A. amurensis</i>		<i>P. pectinifera</i>	
	TC (M)	$E_{max} \pm SEM$ (%)	TC (M)	$E_{max} \pm SEM$ (%)
SALMFa-S1	10^{-8}	19.92 ± 0.99	10^{-8}	14.82 ± 1.38
SALMFa-S2	5×10^{-8}	22.10 ± 2.77	5×10^{-8}	13.61 ± 1.53
SALMFa-A	10^{-8}	11.41 ± 0.61	10^{-7}	8.86 ± 0.90
SALMFa-B	10^{-8}	15.32 ± 0.98	10^{-8}	14.00 ± 1.50

TC is the concentration at which each neuropeptide started to exert myoactivity. The E_{max} is based on the best-fit top value on a dose-dependent curve constructed by the *in vitro* bio-assay and estimated as a relative percentage of the muscle maximum contraction induced by acetylcholine 5×10^{-6} M on apical muscle. TC, threshold concentration; E_{max} , peptide efficacy.

showed a similar pattern of potency. However, the relaxing effect of native SALMFa-A was dramatically lower than those of control SALMFa-S1 and SALMFa-S2 peptides (Fig. 3A). The efficacy of SALMFa-A was roughly 2-fold lower than those of SALMFa-S1 and SALMFa-S2 peptides (Fig. 3B). The pharmacological activity of the SALMFa-S1, SALMFa-S2, SALMFa-A, and SALMFa-B showed $14.82 \pm 1.38\%$, $13.61 \pm 1.53\%$, $8.86 \pm 0.90\%$, $14.00 \pm 1.50\%$, respectively. The TC for SALMFa-S1 and SALMFa-B were observed at 10^{-8} M. On the other hand, the TC of SALMFa-S2 and SALMFa-A was observed at 5×10^{-8} M and 10^{-7} M concentration, respectively (Table 1). The order of relaxing effect on *P. pectinifera* apical muscle was: SALMFa-S1 >




Fig. 3. Investigation of relaxing effects of SALMFamide peptides on *Patiria pectinifera* apical muscle. (A) Concentration dependent response curve of SALMFamide peptides on *P. pectinifera* apical muscle preparation. The sigmoidal-shaped four-parameter dose-response curves were constructed using GraphPad PRISM 8.4. The relaxing response expressed as % of the maximal response to acetylcholine, 5 × 10⁻⁶ M. Data represent the mean ± SD determined from three separate experiments (n = 3). (B) Comparison of the SALMFamide peptides relaxing activity efficacy (E_{max} value). The asterisks on the top of bars denotes statistically significant differences (p < 0.05) determined by one-way analyses of variance followed by the Bonferroni multiple range test.

SALMFa-B > SALMFa-S2 > SALMFa-A.

Discussion

In this study, the SALMFamide peptide isotype and its analog bioactivity was described for the first time. All these obtained results suggest that derivative peptide SALMFa-B has equal or potent relaxing activity compared to native SALMFa-A peptide activity on apical muscle of the starfishes *A. amurensis* and *P. pectinifera*, respectively. The starfish *A. amurensis* is phylogenetically close to *A. rubens* compared to starfish *P. pectinifera* species. Meanwhile, *P. pectinifera* is phylogenetically close to *Patiria miniata* (Fig. 4A). Therefore, the sequences of derivative SALMFa-B peptide were compared with native SALMFa-A peptide of *A. rubens*, and SALMFa peptide of *P. miniata* (Fig. 4B). Firstly, comparison revealed the 66.67% identity between the derivative SALMFa-B and native SALMFa-A peptide of *A. rubens*. The lower similarity of these two peptides may explain the relatively equal lower relaxing effect on *A. amurensis* apical muscle preparation. Secondly, the comparison between the derivative SALMFa-B peptide of *A. rubens* and SALMFa peptide of *P. miniata* showed the 77.78% identity. These relatively higher sequence similarities may elucidate the relatively higher relaxing effect of derivative SALMFa-B of *A. rubens* on apical muscle of *P. pectinifera* compared to those of native SALMFa-A peptide. On the




Fig. 4. Sequence alignment of SALMFamide peptides. (A) Phylogenetic tree constructed by phyloT: phylogenetic three generator. NCBI taxonomy (<https://phylo.t.biobyte.de>). (B) Sequence alignment of derivative SALMFa-B peptide with SALMFamide peptides of *Asterias rubens* and *Patiria miniata*.

other hand, the SALMFa-A and SALMFa-B peptides did not show any activities on tube feet preparation of both starfishes *A. amurensis* and *P. pectinifera* (data not shown). Whereas, the slight relaxation activities were noticed on *P. pectinifera* cardiac stomach preparations at 10⁻⁵ M concentration (Fig. 5). All these results suggest that native SALMFa-A peptides possess main activity on apical muscle that is used by starfish to movement. The derivative SALMFa-B (AYHSALMFamide) peptide with three


P. pectinifera cardiac stomach

Fig. 5. Illustration of relaxing activities on cardiac stomach of starfish *Patiria pectinifera*. The high concentration of SALMFamide (10^{-5} M) peptides applied on cardiac stomach preparation of *P. pectinifera*. The muscle precontractions were caused by carbachol, 5×10^{-6} M.

amino acid substitutions (Ser⁴, Ala⁵, Met⁷) showed relatively high activity than native SALMFa-A (AYHTGLPFamide), suggesting that C-terminal SALMFamide motif has more effectivity on relaxing activity than GLPFamide motif. Also, it seems that the C-terminal region plays a more important role in the receptor binding of starfish smooth muscles than the N-terminal region of SALMFamide peptides. The both SALMFamide-S1 and SALMFamide-S2 showed 100% relaxing response at 10^{-5} M concentration on *A. rubens* cardiac stomach, meanwhile, in this study, SALMFamide-S1 and SALMFamide-S2 did not show significant relaxing response at 10^{-5} M on *P. pectinifera* cardiac stomach (Melarange et al., 1999). These results imply that there are dramatic variances on the SALMFamide's receptor structures in the cardiac stomach of these starfishes. On the other hand, the experiments were carried out under different conditions, which can also affect the results. The relaxing effect of SALMFamide-S1 and SALMFamide-S2 on *A. rubens* apical muscle preparation was determined as minor effect compare to relaxing effect on *A. rubens* cardiac stomach. The relaxation activity of SALMFamide-S1 and SALMFamide-S2 on *A. rubens* apical muscle at 10^{-5} M was 11% and 32%, respectively (Melarange & Elphick, 2003). These results are comparable to the relaxation activity of SALMFamide-S1 and SALMFamide-S2 on *P. pectinifera* and *A. amurensis* apical muscle shown in this study. Therefore, these results suggest that SALMFamide-S1 and SALMFamide-S2 do not show the trivial activity on *A. rubens*, *A. amurensis* and *P. pectinifera* apical muscle preparations.

Competing interests

No potential conflict of interest relevant to this article was re-

ported.

Funding sources

This work was supported by a Research Grant of Pukyong National University in 2021 (C-D-2021-0773).

Acknowledgements

Not applicable.

Availability of data and materials

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Ethics approval and consent to participate

This article does not require IRB/IACUC approval because there are no human and animal participants.

ORCID

Anastasia Kubarova <https://orcid.org/0000-0001-5731-9184>
 Hye-Jin Go <https://orcid.org/0000-0002-5833-4703>
 Hye Young Oh <https://orcid.org/0000-0003-3895-6211>
 Nam Gyu Park <https://orcid.org/0000-0002-7816-1491>

References

- Birenheide R, Tamori M, Motokawa T, Ohtani M, Iwakoshi E, Muneoka Y, et al. Peptides controlling stiffness of connective tissue in sea cucumbers. *Biol Bull.* 1998;194:253-9.
- Byrne M, Mazzone F, Elphick MR, Thorndyke MC, Cisternas P. Expression of the neuropeptide SALMFamide-1 during

- regeneration of the seastar radial nerve cord following arm autotomy. *Proc R Soc B*. 2019;286:20182701.
- Elphick MR. SALMFamide salmagundi: the biology of a neuropeptide family in echinoderms. *Gen Comp Endocrinol*. 2014;205:23-35.
- Elphic MR, Achhala S, Martynyuk N. The evolution and diversity of SALMFamide neuropeptides. *PLOS ONE*. 2013;8:e59076.
- Elphick MR, Emson RH, Thorndyke MC. FMRFamide-like immunoreactivity in the nervous system of the starfish *Asterias rubens*. *Biol Bull*. 1989;177:141-5.
- Elphick MR, Newman SJ, Thorndyke MC. Distribution and action of SALMFamide neuropeptides in the starfish *Asterias rubens*. *J Exp Biol*. 1995;198:2519-25.
- Elphick MR, Price DA, Lee TD, Thorndyke MC. The SALMFamides: a new family of neuropeptides isolated from an echinoderm. *Proc Biol Sci*. 1991;243:121-7.
- Kato S, Tsurumaru S, Taga M, Yamane T, Shibata Y, Ohno K, et al. Neuronal peptides induce oocyte maturation and gamete spawning of sea cucumber, *Apostichopus japonicus*. *Dev Biol*. 2009;326:169-76.
- Kim CH, Kim EJ, Go HJ, Oh HY, Lin M, Elphick MR, et al. Identification of a novel starfish neuropeptide that acts as a muscle relaxant. *J Neurochem*. 2016;137:33-45.
- Klavdieva MM. The history of neuropeptides I. *Front Neuroendocrinol*. 1995;16:293-321.
- Melorange R, Elphick MR. Comparative analysis of nitric oxide and SALMFamide neuropeptides as general muscle relaxants in starfish. *J Exp Biol*. 2003;206:893-9.
- Melorange R, Potton DJ, Thorndyke MC, Elphick MR. SALMFamide neuropeptides cause relaxation and eversion of the cardiac stomach in starfish. *Proc R Soc Lond B*. 1999;266:1785-9.
- Semmens DC, Dane RE, Pancholi MR, Slade SE, Scrivens JH, Elphick MR. Discovery of a novel neurophysin-associated neuropeptide that triggers cardiac stomach contraction and retraction in starfish. *J Exp Biol*. 2013;216:4047-53.
- Semmens DC, Elphick MR. The evolution of neuropeptide signalling: insights from echinoderms. *Brief Funct Genomics*. 2017;16:288-98.
- Thorndyke MC, Carnevali MDC. Regeneration neurohormones and growth factors in echinoderms. *Can J Zool*. 2001;79:1171-208.
- Tinoco AB, Barreiro-Iglesias A, Yañez Guerra LA, Delroisse J, Zhang Y, Gunner EF, et al. Ancient role of sulfakinin/cholecystokinin-type signalling in inhibitory regulation of feeding processes revealed in an echinoderm. *ELife*. 2021;10:e65667.
- Vishwanatha KS, Sobota JA, Eipper BA, Mains RE. Neuropeptide synthesis and storage. In: Stein J, editor. Reference module in neuroscience and biobehavioral psychology. Amsterdam: Elsevier; 2017.