



Using Ivermectin for treating channel catfish (*Ictalurus punctatus*) infected with *Dollfustrema bagarii*

Manh Duc Vu^{1,2}, Kim Minh Anh¹, Lua Thi Dang², Hung Manh Nguyen³, Trinh Tran Thi¹, Ninh Doan Thi¹, Manh Van Ngo⁴, Kim Van Van^{1,*}

¹ Faculty of Fisheries, Vietnam National University of Agriculture, Hanoi 08443, Vietnam

² Research Institute for Aquaculture No. 1, Bac Ninh 16000, Vietnam

³ Institute of Ecology and Biological Resources, Graduate University of Science and Technology, Vietnam Academy of Science and Technology, Hanoi 10072, Vietnam

⁴ Institute of Aquaculture, Nha Trang University, Nha Trang City 650000, Vietnam

Abstract

Channel catfish (*Ictalurus punctatus*), an introduced species in Vietnam, is widely cultured in the Northern provinces. The off-white grub disease in Channel catfish, caused by metacercariae of *Dollfustrema bagarii* (Digenea: Bucephalidae) lodged in internal organs, often results in heavy economic losses. Up to the present, there have been no specific guidelines for preventing and treating this disease. Here, we explore the potential treatment of infected channel catfish through the injection of Ivermectin. We evaluated the tolerance of channel catfish to the drug and determined the optimal dosage for treating off-white grubs disease. Healthy fish weighing 180–200 g received dosages of up to 3.250 mg/kg of body weight. The median lethal dose (LD₅₀) throughout a 24-hour period was 0.808 mg/kg body weight, with a confidence interval ranging from 0.583 to 1.118 mg/kg body weight. The infected fish used for treatment testing ranged in weight from 400 to 500 g. The value for the 24-hour median effective dose (ED₅₀) was 0.253 mg, and the appropriate therapeutic injection ranged from 0.300 to 0.700 mg per kg of body weight. Flowing the histopathological alterations, after the metacercariae were shriveled and died, the immune cells cleaned and eliminated them from fish.

Keywords: *Dollfustrema bagarii*, Effective dose (ED₅₀), *Ictalurus punctatus*, Ivermectin, lethal dose (LD₅₀)

Introduction

In recent years, channel catfish (*Ictalurus punctatus*) has become an important and highly economical species for aquaculture in

the northern region of Vietnam (Hoai et al., 2019; Truong et al., 2020). Diseases in cultured *I. punctatus*, however, have compromised profitability for farmers. Common diseases observed in *I. punctatus* include hemorrhage, ulcers, and both internal and

Received: Mar 20, 2024 Revised: May 16, 2024 Accepted: Jun 11, 2024

*Corresponding author: Kim Van Van

Faculty of Fisheries, Vietnam National University of Agriculture, Hanoi 08443, Vietnam

Tel: +84-904289042, E-mail: kvvan@vnua.edu.vn

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © 2024 The Korean Society of Fisheries and Aquatic Science

external parasitic infections (De la Cruz et al., 2017; Hoai et al., 2019; Wagner et al., 2003). One particularly damaging disease is the “off-white grub” disease, caused by metacercariae of *Dollfus-trema bagarii* (Digenea: Bucephalidae), resulting in substantial economic losses (Kim et al., 2022). Vu et al. (2022) highlighted a staggering 59.7% prevalence of infection in cage cultured channel catfish. Unfortunately, there is currently no effective treatment for this disease in fish.

Ivermectin is widely used to treat parasitic diseases in both humans and terrestrial animals and has shown its effectiveness in treating diseases in aquatic animals as well (Kilmartin et al., 1996; Lorio, 1989; Palmer et al., 1987). Studies have demonstrated its efficacy in treating sea lice on marine fish at three distinct stages through injections (Palmer et al., 1987). Kilmartin et al. (1996) highlighted that salmon is more sensitive to Ivermectin compared to freshwater trout, and both injection and oral administration proved highly effective in treating parasitic diseases. It is worth noting that Ivermectin is a highly toxic drug and can easily kill fish if used in excessive amounts. The signs of Ivermectin poisoning observed in fish include lethargy leading to sinking to the bottom, changes in skin color causing respiratory failure, rapid and erratic swimming, and jumping. However, determining the precise dosage of medication administered via feeding remains a challenge. Therefore, this study focused on evaluating the toxicity of the drug to fish and investigating the optimal dosage for treatment off-white grub disease through injection.

Materials and Methods

Ethics statement

The study was approved by Vietnam National University of Agriculture. All applicable international, national, and institutional guidelines for the care and use of animals were followed.

Experimental preparation and fish injection

Ivermectin concentrations used for injection were determined based on the research findings of Nguyen et al. (2020), when Ivermectin was used to treat an intestinal giant cystic disease caused by *Thelohanellus kitauei* infecting common carp (*Cyprinus carpio*). The varying dose concentrations of ivermectin are detailed in Table 1. The fish were closely monitored for a period of 10 days post injection. The effectiveness of the drug was evaluated based on the prevalence and intensity of infection before and after the experiment. The criteria for determining the infection status included assessing the presence versus absence, as

well as the viability versus mortality, of *D. bagarii* larvae inside the metacercarial cysts in the liver, kidney, spleen, and other organs of the channel catfish.

Before commencing the experiments, we conducted examinations on 3 healthy and 10 infected fish to confirm their infection status and quantify the metacercaria load. The healthy fish were found to be free from metacercariae, while the infected fish exhibited a metacercariae density of 3.37 ± 0.99 metacercariae per gram of liver.

Determining the therapeutic dose

We evenly distributed forty-eight infected fish into six distinct experimental tanks, each containing 500 liters of well water. Based on the lethal dose (LD_{50}) and effective dose (ED_{50}) findings, we determined the range of Ivermectin concentrations used in the injection dose, as outlined in Table 2, to find out the suitable therapeutic dose. Three days after the injection, we examined four fish per tank to assess the intensity of infection in various internal organs and observe any changes in the presence of off-white grub on the surface. On the sixth day post-injection, we examined all the remaining fish.

Calculating fomula

ED_{50} were calculated following the fomula of Kärber (1931).

Statistical analysis

LD_{50} values were determined using probit analysis in STATA/IC 12 (StataCorp, College Station, TX, USA). We conducted an analysis on infection intensity, measured as the number of metacercariae per gram of liver. The metacercariae counts were log-transformed, and we used linear regression with the dose as a continuous variable. Differences with p -values below 0.05

Table 1. Results of detecting the effective dose of ivermectin injection for the treatment of the off-white grub disease in channel catfish

Injection dose (mg Ivermectin/kg body weight)	Number of experimental fish	Number of recovered fish
0	8	0
0.10	8	0
0.20	8	3
0.35	8	6
0.55	8	8
0.70	8	8

Table 2. Efficacy of ivermectin in the treatment of the off-white grub disease in channel catfish after 3 and 6 days of treatment

Injection dose (mg Ivermectin/kg body weight)	Number of experimental fish	3 rd day post-injection (n=4)		6 th day post-injection (n=4)	
		Treatment effect	Infection intensity index (metacercariae/g liver)	Treatment effect	Infection intensity index (metacercariae/g liver)
0.20	8	–	2.61 ± 0.37	–	1.80 ± 0.78
0.30	8	–	1.26 ± 0.26	+	0
0.40	8	+	0	+	0
0.50	8	+	0	+	0
0.60	8	+	0	+	0
0.70	8	+	0	+	0

were considered significant.

Results

Experimented fish and environmental characteristics

Throughout the trial period, the temperature fluctuated between 28 °C to 35 °C, with an average of 32.67 °C. Dissolved oxygen levels varied from 3.5 to 7.5 mg/L, with an average concentration of 5.5 mg/L. The pH values were observed within the range of 6.8 to 7.7, with an average pH of 7.3. Additionally, NH₃⁺ levels ranged from 0 to 1 mg/L, averaging at 0.35 mg/L, while NO₂⁻ levels ranged from 0 to 0.5 mg/L, with an average concentration of 0.15 mg/L.

Toxicity of ivermectin on channel catfish

Most poisoned fish exhibited immediate symptoms and died within 2–22 hours, typically within 4–10 hours post injection. Specifically, injecting 2.75 mg of Ivermectin per kg of body weight resulted in the mortality of all fish within 24 hours post-injection. Those fish that survived beyond the initial 24 hours period remained alive until the conclusion of the 96-hour experiment. The LD_{50-24h} value was determined to be 0.808 mg/kg body weight, with a confidence interval ranging from 0.583 to 1.118 mg/kg body weight (Table 3).

Treatment dose of ivermectin for the off-white grub disease in channel catfish

The detection of effective dose (ED₅₀)

No abnormal behavior was observed in experimental fish at the onset of testing the effective dose at 0.10 mg of Ivermectin/kg body weight (Table 1). As the ED₁₀₀ was determined to be 0.55 mg of Ivermectin per kg of body weight, the calculated value for

Table 3. Ivermectin concentrations used in injection dose to determine the toxicity (lethal dose, LD₅₀) of channel catfish

Injection dose (mg Ivermectin/kg body weight)	Number of experimental fish	Number of death fish	Death rate
0	4	0	0/4
0.5	4	0	0/4
0.75	4	1	1/4
1.25	4	2	2/4
1.75	4	3	3/4
2.25	4	3	3/4
2.75	4	4	4/4
3.25	4	4	4/4

ED₅₀ was to be 0.253 mg/kg body weight.

At the dose of 0.70 mg Ivermectin per kg of body weight, no fish experienced mortality. However, additional toxic effects were observed, including decreased feeding behavior, lethargy, disorientation, and sinking to the bottom within the initial few hours.

Results of determining the therapeutic dose

The evaluation of the therapeutic dose ranged from 0.20 to 0.70 mg Ivermectin per kg of body weight. Even after administering an injection of 0.20 mg Ivermectin per kg of body weight, live *D. bagarii* metacercariae were still detected in the fish, with infection intensities of 2.61 ± 0.37 and 1.80 ± 0.78 metacercariae per gram of liver after 3 and 6 days of treatment, respectively.

After three days of treatment with an injection dose of 0.30 mg Ivermectin per kg of body weight, the infection intensity decreased to 1.26 ± 0.26 metacercariae per gram of liver. By sixth day of observation, no *D. bagarii* metacercariae were found (Fig. 1). Similarly, injections of 0.40, 0.50, 0.60, and 0.70 mg/kg of




Fig. 1. Liver of channel catfish. (A) with dense of metacercariae (black arrows) before treatment; (B) metacercariae were exterminated after treatment (black cycles) six days.

body weight with Ivermectin produced positive effect (Table 2).

Histopathological alterations in liver and kidney tissue before and after injection

Prior to treatment, metacercariae infesting the liver and kidneys manifested as gross lesions (Fig. 2A–2D). Metacercariae evaded cysts (Fig. 2A and 2B) and infiltrated the tissue, resulting in regions of tissue exhibiting lax structure and hemorrhaging. After six days of treatment, the cysts and metacercariae underwent organ-atrophy, accompanied by the emergence of numerous pigmented macrophages tasked with clearance (Fig. 2E and 2F). The cysts disengaged from liver and kidney, resulting in the formation of concave scar areas (Fig. 2G and 2H).

Discussion

The range of environmental parameters experienced during these trials, as highlighted by Alejandro Buentello et al. (2000), Boyd & Pillai (1985), and Boyd et al. (2002), was suitable for the development of channel catfish.

In aquatic animals, the detection of medical toxicity usually occurs rapidly through observable signs. These signs include restlessness, frantic movement, gulping of air, accumulation of mucus on the body, loss of equilibrium by swimming sideways,

and eventual collapse leading to the demise of the fish (Andem et al., 2015). In instances of Ivermectin poisoning due to oral administration in small mammals like mice and rats, symptoms observed encompass vomiting, central nervous system depression, acute respiratory distress, and dilated pupils (Trailović & Varagić, 2007). Throughout our study, fish exhibited poisoning symptoms such as sinking to the bottom, alterations in skin tone, rapid and erratic swimming, and jumping. No additional signs of illness or infection were noted, and notably, all fish swiftly recuperated following the trial. Consequently, based on these observations, it can be reasonably inferred that Ivermectin poisoning served as the primary cause of fish mortality.

Despite numerous studies on Ivermectin's toxicity to the environment and other organisms, limited data exists regarding the LD₅₀ of Ivermectin injection in channel catfish. Reports on the ecotoxicity of Ivermectin across different aquatic species reveal varying levels: 0.003 ppm for rainbow trout (*Oncorhynchus mykiss*), 0.048 mg/L for 96 hours in bluegill sunfish (*Lepomis macrochirus*), 2.5×10^5 mg/L for 48 hours in water flea (*Daphnia magna*), and over 9.1 mg/L for 72 hours in green algae (*Pseudokirchneriella subcapitata*) (MSD, 2022; OECD, 2011). It is important to note that LC₅₀ or LD₅₀ doses may differ based on species, drug administration routes, and specific time points (MSD, 2022). Ivermectin is categorized as highly toxic accord-




Fig. 2. Histopathological alterations in liver and kidney tissue (H&E stained sections) before (A–D) and after (E–H) six days of injection. Metacercariae (black arrows) before (A: liver, B: kidney) treating and the later (E, F: liver); Structure-loose (C: liver) and degeneration (B, D: kidney) were observed in fish internal organs; Immune cells become active around damaged areas (white arrows – eosinophilic granular cells and yellow arrows – pigmented macrophages); Scars in liver (G, H) three-day-post treating (black cycles).

ing to the Hazard Classification System - GHS (United Nations, 1992). However, this classification markedly differs concerning terrestrial animals, particularly mammalian species. The LD₅₀ of

Ivermectin in rats, mice, and monkeys after oral administration (a single dosage) is approximately 10–50 mg/kg, 25 mg/kg, and 24 mg/kg body weight, respectively. For instance, the LC_{50-24h} by

skin contact for rabbits was 406 mg Ivermectin/kg body weight, and LC_{50-24h} for rats was over 660 mg/kg body weight. Ivermectin adversely affects the respiratory system of rats (LC_{50-24h} was 5.11 mg/L of breathing air). In mice, the LD_{50} via oral administration was found to be 18 mg Ivermectin/kg body weight (Trai-ović & Varagić, 2007), whereas in beagle dogs, it was 80 mg/kg body weight (Hopper et al., 2002). From this data, it is evident that aquatic animals demonstrate sensitivity to Ivermectin toxicity compared to mammalian species. Additionally, their sizes play a role in their tolerance to Ivermectin toxicity. Therefore, further detailed studies on Ivermectin toxicity in channel catfish are warranted, considering varied sizes and smaller differences between consecutive injection doses.

Beside highly toxic, Ivermectin also caused residues in food products (Qiao et al., 2023). In 70 days Wang et al. (2020) observed, the concentration of Ivermectin reached a peak in the muscle (0.1921 $\mu\text{g/g}$) and visceral mass (0.25982 $\mu\text{g/g}$) of brocarded carp at 7 days after immersing in the water at a concentration of 0.16 $\mu\text{g/L}$, and then gradually decreasing to 10.179 ng/g in the visceral mass and 3.823 ng/g in the muscle. After administered Ivermectin at the dose level of 0.1 mg/kg bw, the highest total radioactive residue concentrations of 53, 45, and 44 ng/g were obtained on postdose day 1 for hybrid striped bass, largemouth bass, and yellow perch, respectively (Shaikh et al., 2012). Normally, treatment of aquatic animals in large numbers often uses the oral administration. However, there have been no studies to determine residues on channel catfish muscle after injection. Meanwhile, the research of Edwards et al. (1988) in 12 healthy male resulted that after a single dose of 12 mg/kg bw follow oral administration, the maximum plasma concentration reached the highest point at 880 ng/h/mL without any clinical side effects. Those values were lower than the acute oral toxicity in human, at 1,511 mg/kg bw (MSD, 2022). However, the assessment of the effects of long-term absorption needs to be clarified.

Ivermectin demonstrated effective control of “yellow spot” disease caused by *Clinostomum marginatum* at a dose of 0.22 mg/kg body weight (Lorio, 1989). Similarly, successful treatment of intestinal cysts induced by *T. kitauei* involved oral administration of 0.6 mg of Ivermectin per kg of body weight for three consecutive days (Nguyen et al., 2020). However, studies investigating its applications in channel catfish have been limited. Further research in this subject is essential to enhance the practical utilization of this medication.

In conclusion, Ivermectin exhibited high toxicity towards

channel catfish. An injection dose of 1.406 mg per kg of body weight resulted in a 50% mortality rate among 180–200 g channel catfish within 24 hours. Effective treatment for the off-white grub disease caused by metacercariae of *D. bagarii* required a dose of 0.2525 mg/kg body weight. Doses ranging from 0.3 to 0.7 mg/kg body weight of Ivermectin successfully treated diseases caused by *D. bagarii* metacercariae in channel catfish internal organs. Notably, Ivermectin emerged as a valuable treatment for the off white grub disease caused by *D. bagarii* metacercariae in channel catfish. For practical application in treating large numbers of fish in cages, more research on oral 182 administration is needed.

Competing interests

No potential conflict of interest relevant to this article was reported.

Funding sources

This work was supported by VNUA (Vietnam National University of Agriculture) research project no T2023-14-52. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

I am very grateful to Hoa Dang for assistance with data gathering, and to Henry Madsen for assistance with data processing and constructive comments on an earlier version of the manuscript.

Availability of data and materials

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Ethics approval and consent to participate

The study was approved by Vietnam National University of Agriculture. All applicable international, national, and institutional guidelines for the care and use of animals were followed.

ORCID

Manh Vu Duc	https://orcid.org/0009-0003-6025-0159
Kim Minh Anh	https://orcid.org/0000-0001-8427-7274
Lua Thi Dang	https://orcid.org/0000-0002-7448-172X
Hung Manh Nguyen	https://orcid.org/0000-0001-5741-419X
Van Manh Ngo	https://orcid.org/0009-0006-1341-2938
Van Van Kim	https://orcid.org/0000-0002-9530-4829

References

- Alejandro Buentello JA, Gatlin DM 3rd, Neill WH. Effects of water temperature and dissolved oxygen on daily feed consumption, feed utilization and growth of channel catfish (*Ictalurus punctatus*). *Aquaculture*. 2000;182:339-52.
- Andem AB, Esenowo IK, Ibor OR, Abosi A. Toxicological effects of formaldehyde concentrations on African Cat fish, *Clarias gariepinus* fingerlings. *Int J Fish Aquat Stud*. 2015;5:75-9.
- Boyd CE, Pillai VK. Water quality management in aquaculture. Kochi: CMFRI Special Publication; 1985.
- Boyd CE, Wood CW, Thunjai T. Aquaculture pond bottom soil quality management [Pond Dynamics/Aquaculture Collaborative Research Support Program]. Corvallis, OR: Oregon State University; 2002.
- De la Cruz NI, Rabago JL, Monreal AE, Colín VH, Aguirre G, Merino JO, et al. Diagnosis and frequency of parasites in channel cat fish (*Ictalurus punctatus*) on northeastern and gulf coast farms Mexico. *Int J Fish Aquat Stud*. 2017;5:98-103.
- Edwards G, Dingsdale A, Helsby N, Orme MLE, Breckenridge AM. The relative systemic availability of ivermectin after administration as capsule, tablet, and oral solution. *Eur J Clin Pharmacol*. 1988;35:681-4.
- Hoai TD, Trang TT, Van Tuyen N, Giang NTH, Van Van K. *Aeromonas veronii* caused disease and mortality in channel catfish in Vietnam. *Aquaculture*. 2019;513:734425.
- Hopper K, Aldrich J, Haskins SC. Ivermectin toxicity in 17 col-lies. *J Vet Intern Med*. 2002;16:89-94.
- Kärber G. Beitrag zur kollektiven Behandlung pharmakologischer Reihenversuche. *Naunyn-Schmiedebergs Arch Exp Pathol Pharmacol*. 1931;162:480-3.
- Kilmartin J, Caxabon D, Smith P. Investigations of the toxicity of the ivermectin for salmonids. *Bull Eur Assoc Fish Path*. 1996;17:58-4.
- Kim VV, Nguyen MH, Greiman SE, Nguyen HV, Nguyen CN, Vu MD, et al. Molecular and morphological characterization of *Dollfustrema bagarii* (Digenea: Bucephalidae) metacercariae from aquaculture channel catfish (*Ictalurus punctatus*) in northern Vietnam. *J Fish Dis*. 2022;45:1165-71.
- Lorio WJ. Experimental control of metacercariae of the yellow grub *Clinostomum marginatum* in channel catfish. *J Aquat Anim Health*. 1989;1:269-71.
- Merck Sharp & Dohme [MSD]. Ivermectin (3.5%) formulation [Internet]. 2022 [cited 2024 Feb 4]. [https://www.msd.com/docs/product/safety-data-sheets/ah-sds/Ivermectin%20\(3.5_pct\)%20Formulation_AH_JP_6N.pdf](https://www.msd.com/docs/product/safety-data-sheets/ah-sds/Ivermectin%20(3.5_pct)%20Formulation_AH_JP_6N.pdf)
- Nguyen VT, Tran TD, Pham TT, Doan TN, Kim VV, Truong DH. Experimental treatment of intestinal giant cystic disease caused by *Thelohanellus kitauei* infecting common carp (*Cyprinus carpio*). *Vietnam J Agric Sci*. 2020;18:1139-9.
- Organization for Economic Cooperation and Development [OECD]. Test no. 201: freshwater alga and cyanobacteria, growth inhibition test, OECD guidelines for the testing of chemicals, section 2 [Internet]. OECD, Paris. 2011 [cited 2024 Feb 4]. <https://doi.org/10.1787/9789264069923-en>
- Palmer R, Rodger H, Drinan E, Dwyer C, Smith PR. Preliminary trials on the efficacy of ivermectin against the parasitic copepods of Atlantic salmon. *Bull Eur Assoc Fish Pathol*. 1987;7:47-54.
- Qiao L, Xu J, Yang Z, Li X, Chen L, Sun H, et al. Residual risk of avermectins in food products of animal origin and their research progress on toxicity and determination. *Food Rev Int*. 2023;39:7019-47.
- Shaikh M, Valmaggia L, Broome MR, Dutt A, Lappin J, Day F, et al. Reduced mismatch negativity predates the onset of psychosis. *Schizophr Res*. 2012;134:42-8.
- Trailović SM, Varagić VM. The effect of ivermectin on convulsions in rats produced by lidocaine and strychnine. *Vet Res Commun*. 2007;31:863-72.
- Truong DH, Kim VV, Dao LA, Nguyen TH, Nguyen VT, Vu DM, et al. Pathological characterization and PCR application for diagnosis *Edwardsiella ictaluri* infecting Channel catfish (*Ictalurus punctatus*). *Vietnam J Agric Sci*. 2020;18:94-11.
- United Nations. Globally harmonized system of classification and labeling of chemicals (GHS). In: Proceedings of the Rio Conference on Environment and Development; 1992; Rio de Janeiro, Brazil.
- Vu DM, Kim MA, Nguyen MH, Do DH, Truong DH, Dang TL, et al. White spot disease on internal organs of channel catfish (*Ictalurus punctatus*) cage culture in northern Vietnam caused by metacercariae of *Dollfustrema bagarii*. *Vietnam J Vet Sci Technol*. 2022;29:62-8.
- Wagner B. Catfish 2003, part II: reference of foodsize catfish health and production practices in the United States (No. 1499-2016-130721). Fort Collins, CO: National Animal

Health Monitoring System; 2003.

Wang D, Han B, Li S, Du X, Cao Y, Lu T. Assessment of the fate and effect of ivermectin in a simulated aquaculture ecosystem. *Aquac Res* 2020;51:535-41.